

**Results of ALPS Treated Water Marine Monitoring:
Marine biota survey (carbon-14 in fish) (October 2023)**

1. Outline of survey

(1) Date of sampling

October 18, 2023

(2) Sampling points

3 sampling points on coastal waters in the Fukushima Prefecture

(3) Detail of the survey

- The measurements of radioactive material concentration (carbon-14) in marine biota (fish)

Analysis with target lower limit of detection of 2 Bq/kg-fresh.

*A target lower limit of detection means a value that is set for quality control to assure at least the detection up to the value when analysis is conducted. Each actual lower limit of detection differs according to samples, and is equal to or lower than a target lower limit of detection.

2. Outline of results

(1) Marine biota survey (3 sampling points (8 samples))

Concentrations of carbon-14 in the marine biota (fish) range from 19 Bq/kg-fresh to 29 Bq/kg-fresh.

* In this survey, only two samples (usually three) were collected at some of the stations with small quantities, and some of the samples collected in small quantities were mixed with multiple fish species (usually one fish species per sample).

These results were approximately equal to results of carbon-14 analysis in marine biota (fish) conducted in past surveys.

The range of carbon-14 specific radioactivity of marine biota in this survey:

230 Bq/kg-carbon to 240 Bq/kg-carbon (19 Bq/kg-fresh to 29 Bq/kg-fresh)

The range of carbon-14 specific radioactivity of marine biota (fish) in past surveys:

230 Bq/kg-carbon to 250 Bq/kg-carbon (16 Bq/kg-fresh to 28 Bq/kg-fresh)

* A specific radioactivity means the radioactivity per unit mass of a substance containing radioactive isotopes. In the case of Bq/kg-carbon above, it represents the radioactivity per 1 kg of carbon in the sample. It is difficult to compare radioactivity concentrations due to differences in the carbon content rate in samples of marine biota, specific radioactivity is also shown as reference information.

(Detailed are attached)

(Maps attached)

Attachment

Analysis results for carbon-14 in marine biota (fish)

Sampling point	Sampling date (yyyy/mm/dd)	Species	Sampling depth (m)	Nuclide	Radioactivity concentration ^{*1,*2}	Unit
E-SF1	2023/10/18	Mixed fishes A	-	C-14	25 ± 0.4	Bq/kg-fresh
E-SF1	2023/10/18	Mixed fishes B	-	C-14	26 ± 0.4	Bq/kg-fresh
E-SF2	2023/10/18	<i>Paralichthys olivaceus</i>	-	C-14	26 ± 0.4	Bq/kg-fresh
E-SF2	2023/10/18	<i>Pagrus major</i>	-	C-14	29 ± 0.4	Bq/kg-fresh
E-SF2	2023/10/18	<i>Triakis scyllium</i>	-	C-14	24 ± 0.4	Bq/kg-fresh
E-SF3	2023/10/18	<i>Paralichthys olivaceus</i>	-	C-14	23 ± 0.3	Bq/kg-fresh
E-SF3	2023/10/18	<i>Okamejei schmidtii</i>	-	C-14	19 ± 0.3	Bq/kg-fresh
E-SF3	2023/10/18	<i>Squatina japonica</i>	-	C-14	22 ± 0.3	Bq/kg-fresh

*1 Radioactivity concentrations are presented as radioactivity concentration ± combined standard uncertainty.

*2 Values below detection limit are shown by lower limit of detection (e.g., “<10 Bq/ kg-fresh” indicates a value lower than 10 Bq/ kg-fresh).

(Attachment)




Fig. 1 Sampling points of marine biota (fish and seaweed)